

# A polynomial endomorphism of $\mathbb{C}^2$ with a wandering Fatou component

M. Astorg<sup>1</sup>   X. Buff<sup>1</sup>   R. Dujardin<sup>2</sup>   H. Peters<sup>3</sup>   J. Raissy<sup>1</sup>

<sup>1</sup>Université de Toulouse

<sup>2</sup>Université Paris Est de Marne-la-Vallée

<sup>3</sup>Université d'Amsterdam

March 11, 2015

- 1 Introduction
- 2 Strategy of the construction
- 3 Parabolic implosion


## General setting

Let  $M$  be a complex manifold, and  $f : M \rightarrow M$  a holomorphic map.

## Definition

- The Fatou set is the largest open set on which the iterates  $\{f^n, n \in \mathbb{N}\}$  form a normal family.
- The Julia set is the complement of the Fatou set.
- A connected component of the Fatou set is called a Fatou component.

# Example : Julia and Fatou sets of $z \mapsto z^2 - 1$



## Theorem (Sullivan, 1985)

Let  $f : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  be a rational map of degree  $d \geq 2$ . Every Fatou component of  $f$  is preperiodic.

## Consequence :

Together with the classification theorem, we get a complete description of the dynamics in the Fatou set.


## More generally :

If  $f : M \rightarrow M$  is holomorphic, and  $M$  is a complex manifold, must every Fatou component be preperiodic ?

## Answer :

- $M = \mathbb{C}$  and  $f$  a transcendental map : no (Baker, 1976)
- $M = \mathbb{P}^1(\mathbb{C})$ : yes (Sullivan, 1985)
- $M = \mathbb{C}$  and  $f$  transcendental with only finitely many singular values : yes (Eremenko-Lyubich 1992, Goldberg-Keen 1986)
- $M = \mathbb{C}^2$ ,  $f$  biholomorphic and transcendental : no (Fornaess-Sibony, 1998)
- $M = \mathbb{C}$  and  $f$  transcendental with bounded set of singular values : no (Bishop, 2012)
- $M = \mathbb{P}^2(\mathbb{C})$ ,  $f$  polynomial : no (A-Buff-Dujardin-Peters-Raissy, 2014).

A transcendental example :  $z \mapsto z - \sin(z) + 2\pi$



## Theorem (A.-Buff-Dujardin-Peters-Raissy)

There exists a polynomial map  $P : \mathbb{C}^2 \rightarrow \mathbb{C}^2$  with a wandering Fatou component.

## More precisely :


There exists polynomials  $f$  of the form  $f(z) = z + z^2 + O(z^3)$  such that for any polynomial  $g$  of the form  $g(w) = w - w^2 + O(w^3)$ , the skew-product

$$P(z, w) = \left( f(z) + \frac{\pi^2}{4} w, g(w) \right)$$


has a wandering domain, that accumulates the complex line  $\{w = 0\}$ .



# Sketch of the dynamics of $P(z, w) = (f(z) + \frac{\pi^2}{4}w, g(w))$



# Projection of a wandering orbit on $\{w = 0\}$



# Strategy of the construction (idea : M. Lyubich)

Property wanted for  $P(z, w) = (f(z) + \frac{\pi^2}{4}w, g(w))$

We want to chose  $f$  so that there is a subsequence  $m_k$  and an open set  $U$  such that for all  $(z, w) \in U$ ,


$$P^{m_k}(z, w) \rightarrow (\hat{z}, 0)$$

and  $(\hat{z}, 0)$  is not preperiodic for  $P$ .

## Proposition

If  $P$  satisfies that property, then  $P$  has a wandering domain.

# Sketch



# Sketch of the proof

Let  $U$  be a connected open set such that  $\forall (z, w) \in U, P^{m_k}(z, w) \rightarrow (\hat{z}, 0)$   
(\*)

- $U$  is a subset of a Fatou component  $\Omega_0$
- We have (\*) on all of  $\Omega_0$  by analytic continuation
- If  $\Omega_i = P^i(\Omega_0)$ , then for all  $(z, w) \in \Omega_i$ ,  $P^{m_k}(z, w)$  converges to  $(f^i(\hat{z}), 0)$
- If  $i \neq j$ , we have  $f^i(\hat{z}) \neq f^j(\hat{z})$
- Therefore if  $i \neq j$ ,  $\Omega_i \neq \Omega_j$  and so  $(\Omega_n)_{n \in \mathbb{N}}$  is wandering.

# Dynamics near $\{w = 0\}$

## Notation

Let  $f_w(z) = f(z) + \frac{\pi^2}{4}w$ . Then :

$$P^n(z, w_0) = (f_{w_{n-1}} \circ \dots \circ f_{w_0}(z), g^n(w_0))$$

with  $w_k = g^k(w_0) \sim \frac{1}{k}$ .

## Idea

Understanding the dynamics of  $P$  near  $\{w = 0\}$  amounts to understanding the non-autonomous compositions  $f_{w_n} \circ \dots \circ f_{w_k}$ , with  $f_{w_i}(z) \simeq f(z)$

## Notations

- $f(z) = z + z^2 + O(z^3)$  has a parabolic fixed point at 0
- $\mathcal{B}_f$  is its parabolic basin
- $\phi_f$  is the (normalized) attracting Fatou coordinate


$$\phi_f \circ f = \phi_f + 1$$

- $\psi_f$  is the (normalized) repelling Fatou parametrization

$$f \circ \psi_f(Z) = \psi_f(Z + 1)$$

- $L_f = \psi_f \circ \phi_f$  is the (phase 0) Lavaurs map

# Lavaurs map





## Theorem (Lavaurs)

Let  $\epsilon_k \rightarrow 0$  and  $n_k \rightarrow +\infty$ , such that  $\frac{\pi}{\sqrt{\epsilon_k}} - n_k \rightarrow 0$ . Then  $(f + \epsilon_k)^{\circ n_k}$  converges to the Lavaurs map  $L_f$  uniformly on compacts of  $\mathcal{B}_f$ .


## Consequence

With  $\epsilon_k = \frac{\pi^2}{4} w_{k^2} \sim \frac{\pi^2}{4k^2}$  and  $n_k = 2k$ , the sequence  $\underbrace{f_{w_{k^2}} \circ \dots \circ f_{w_{k^2}}}_{2k}$  converges locally uniformly to  $L_f$  (with local uniformity in  $w$  as well).

## Key proposition (A.-Buff-Dujardin-Peters-Raissy)

For all  $w \in \mathcal{B}_g$ , the sequence  $\underbrace{f_{w_{(k+1)^2-1}} \circ \dots \circ f_{w_{k^2}}}_{2k+1}$  converges locally uniformly to  $L_f$  (with local uniformity in  $w$  as well).

# Lavaurs' theorem



## Theorem (A-Buff-Dujardin-Peters-Raissy, 2014)


Suppose  $f(z) = z + z^2 + O(z^3)$  is such that  $L_f$  has an attracting fixed point. Let  $g(w) = w - w^2 + O(w^3)$ . Then

$$P(z, w) = \left( f(z) + \frac{\pi^2}{4} w, g(w) \right)$$

has a wandering domain.


## Numerical example :

$$P(z, w) = \left( z + z^2 + 0.95z^3 + \frac{\pi^2}{4} w, w - w^2 \right)$$




Red dots : attracting fixed points of  $L_f$

Grey part : points that do not escape under  $\langle f, L_f \rangle$  (Julia-Lavaurs set of  $f$ )



# Idea of the proof of the key proposition



$$\text{Notation : } P^k(z, g^{n^2}(w)) = \left( z_k, g^{n^2+k}(w) \right)$$

Thank you for your attention!